EN AS EN Common Name: Cellery

Scientific Name: Apium graveolens

Umbelliferae Family:

## **Recommended Varieties**

Green giant

Utah green

# **Expected Yield**


8 - 10 tonnes/ha

# **Soil Requirement**

Needs a well drained soil with sufficient moisture

Grows well under sarlon shed

Needs an optimum temperature of 6 - 21°C for germination



# **Planting Time**

Grows well during cooler season (April – August)

## **Seed Rate**

250g/ha

## **Spacing**

60cm to 90cm between rows

15 – 25cm between plants

## Land Preparation

Two ploughing and two harrowing is recommended for good soil tilth.

#### Method of Planting

Direct sowing on well prepared beds with fine soil

#### **Fertilizer Requirement**

Poultry manure: 5t/ha broadcast, mix well into soil 2 weeks before planting.

N.P.K: 200kg/ha basal application during planting.

Urea: 100kg/ha – top dress 2 weeks after sowing.

## **Plant protection**

Hand weed regularly or use hoe.

#### **Disease Control**

Anthracnose – control using Anatracol (30g/15 litres of water) or Mancozeb at the rate of 50g/15 litres of water.

#### Harvesting

Harvest within 80 to 110 days (14 - 18 weeks) and get \$24,000 per hectare at \$3.00/kg.



# 

## 13. Food Value


Celery is a good source of Protein, Iron, Vitamin A & C, Thiamin, Riboflavin, Niacin, Potassium, Calcium and Magnesium.





For more information contact: Ministry of Agriculture, Private Mail, Raiwaqa.
Phone: (+679) 338 4233 Fax (+679) 338 7157 / Website: www.agriculture.gov.fj
Facebook: Ministry of Agriculture Fiji Twitter: Fiji Agriculture
Email: agrihelp@govnet.gov.fj





Farmer's Leaflet 2015